
iKnowMedSM Generation 2

## PIMSH17: Oncology: Utilization of Prophylactic GCSF for Cancer Patients Receiving Low-Risk Chemotherapy

In this lesson, you will learn the required documentation for the MIPS# PIMSH17: Oncology: Utilization of Prophylactic GCSF for Cancer Patients Receiving Low-Risk Chemotherapy quality measure. This is an inverse measure.


Patients will be included in the denominator if they have an E&M visit in the reporting period and have an order for GCSF by their established clincian within 7 days of IV or oral chemotherapy.

Established Clinician = 1 or more visits with the patient in the reporting year.




- 1. To add a problem, click on the Problems tab.
- 2. Click on ADD PROBLEM.

## iKnowMedSM Generation 2



- 3. Type the problem in the search box.
- 4. Select the appropriate problem from the list.
- 5. Add the date of diagnosis.
- 6. Identify the problem as active.



- 7. Click on the Details box.
- 8. After details documented, click on SAVE & CLOSE.





## iKnowMedSM Generation 2

- To see if GCSF was ordered within 7 days of chemotherapy, click the Flowsheet tab.
- 10. Check to see if the patient was given IV chemotherapy, indicated by a next to the medication, or if an oral chemotherapy medication was ordered.
- 11. Check to see if a GCSF was ordered within 7 days of the chemotherapy. If so, this will meet the numerator requirement.

This concludes the lesson for the MIPS# PIMSH17: Oncology: Utilization of Prophylactic GCSF for Cancer Patients Receiving Low-Risk Chemotherapy quality measure.